Exposure of Impacted Upper Canine



This leaflet has been designed to improve your understanding of any forthcoming treatment and contains answers to many of the commonly asked questions. If you have any other questions that the leaflet does not answer or would like further explanation please ask.



The problem

The canine, or ”eye tooth”, normally erupts into the mouth between the ages of 11 and 13. Sometimes one or both canines develop in the wrong position. Often they lie across the roof of the mouth behind the front teeth.

Why do I need treatment?

Because one or both of your canines are in the wrong place, as part of your on-going orthodontic treatment it is necessary to help the tooth erupt into the mouth. If left alone the tooth will not erupt normally and may either damage the roots of the front teeth or push them out of position.

What does the treatment involve?

Helping the tooth erupt into your mouth involves a relatively minor surgical procedure. This usually takes place under a “day case” general anaesthetic. This means that although you are put to sleep completely you will be able to go home on the same day as surgery.

How will the orthodontist pull the tooth into the correct position?

Once the canine is exposed one of three things will happen under the same anaesthetic. What is going to happen for you will already have been discussed.

- **Bracket and chain**
  A small bracket is glued to the tooth. Attached to this is a chain which your orthodontist can then use to pull the tooth into the right position. The chain is usually stitched out of the way but it is quite delicate and therefore it is important to be careful when eating for the first few weeks after surgery.

- **A plate**
  Sometimes a small window will be cut in the gum over the tooth and a plastic "dressing" plate put in place to cover the area. This plate is held in your mouth with clips that attach to some of your back teeth. It is important that you wear the plate all the time except when you take it out to clean your teeth. Without the plate the gum may grow back making it difficult for the orthodontist to move the tooth into position. The plate should be kept in mouth for 24 hours after surgery before taking it out to clean. The plate should be worn for 1 week post operatively.

- **A pack**
  Sometimes a pack made from gauze soaked in an antiseptic is placed over the tooth after it is exposed. The pack is kept in position with stitches and removed after a week. You must be careful not to dislodge the pack. If this happens you should contact the department for advice. Sometimes it is necessary to hold the gum back in the right position with stitches at the end of the operation. These are usually dissolvable and take about two weeks to disappear.

Is there much pain or swelling?

Although the procedure isn't particularly painful, you may experience some soreness afterwards. There is usually very little in the way of swelling. If it is likely to be sore your surgeon will arrange painkillers for you. It is not usually necessary to take antibiotics.

Will I need another appointment?

You will need to return 1 week after surgery to have the area checked by your orthodontist.

Further appointments after this will obviously be necessary for your on-going orthodontic care.

This leaflet is specific to the practice of Oral and Maxillo-Facial Surgery in the United Kingdom and is in addition to the generic guidance given in the GMC publications of Good Medical and Surgical Practice 2001 & 2002

If you have any comments about this leaflet or the service you have received you can contact :

Maxillo-Facial Unit
Huddersfield Royal Infirmary
Telephone No: (01484) 342336 / 345486

or

Children's Ward / Ward 4
Telephone No: (01422) 223277 / (01484) 342482

www.cht.nhs.uk

If you would like this information in another format or language contact the above.

Potřebujete-li tyto informace v jiném formátu nebo jazyce, obraťte se prosím na výše uvedené oddělení

Jeżeli są Państwo zainteresowani otrzymaniem tych informacji w innym formacie lub wersji językowej, prosimy skontaktować się z nami, korzystając z ww. danych kontaktowych

ਜੇ ਤੁਸੀਂ ਇਸ ਜਾਣਕਾਰੀ ਬਿਸੇ ਕੋਜ ਪ੍ਰਾਰੂਪ ਜਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਲੈਣਾ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਬਿਜਪਾ ਕਰਕੇ ਉਪਰੋਕਤ ਵਿਭਾਗ ਵਿੱਚ ਸਾਡੇ ਨਾਲ ਸੰਪਰਕ ਕਰੋ।

اگر آپ کو یہ معلومات کسی اور فارمیٹ یا زبان میں درکار ہوں، تو برائے مہربانی مندرجہ بالا شعبے میں ہم سے رابطہ کریں.

"إذا احتجت الحصول على هذه المعلومة بشكل مغاير أو مترجمة إلى لغة مختلفة فيرجى منك الاتصال بالقسم المذكور أعلاه"