Mid Trimester Medical Abortion
13 – 17 + 6 Weeks

Risks And Complications

Immediate complications

- Incomplete abortion
  Retained products/products requiring surgical intervention 90 in 1000 (9%)
- Bleeding
  About 6 in 1000 (0.6%) will require a blood transfusion
- Continuing pregnancy - fetus not expelled after repeated drug regimens 8 in 1000 (0.8%)
- Uterine rupture – 2 in 100,000 cases
- Infection 5 in 100 (5%)

Delayed complications

In the Early Weeks

a) Uterine/pelvic infection – less than 5 in 100

b) Short term emotional distress

compassionate care

Long Term

a) Future reproductive outocome

No proven association between abortion and infertility or ectopic pregnancy. However, it may be associated with a small increase risk of subsequent miscarriage or preterm delivery.

b) Psychological sequalae

These are rare. Some studies suggest a small increase risk of suicide, attempted suicide or self harm after abortion. These may reflect the pre-existing condition.

Breast Cancer Risk

There is no association between induced abortion and breast cancer.

If you have any comments about this leaflet or the service you have received you can contact :

Secretary to Consultant Gynaecologist
Heatherdale Clinic / Gynae Assessment Unit
Calderdale Royal Hospital

Telephone No: 01422 224979 / 224415

www.cht.nhs.uk

If you would like this information in another format or language contact the above.

Potřebujete-li tyto informace v jiném formátu nebo jazyce, obraťte se prosím na výše uvedené oddělení

Jeżeli są Państwo zainteresowani otrzymaniem tych informacji w innym formacie lub wersji językowej, prosimy skontaktować się z nami, korzystając z ww. danych kontaktowych

ਜੇ ਤੁਸੀਂ ਇਸ ਜਾਣਕਾਰੀ ਬਿਸੇ ਹੋਰ ਪਾਰੂਪ ਜਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਲੈਣਾ ਚਾਹੁੰਦੇ ਹੋ, ਤਾਂ ਬਿਚਪਾ ਜਰਰੇ ਉਪਰੋਕਤ ਵਿਭਾਗ ਵਿੱਚ ਸਾਡੇ ਨਾਲ ਸੰਪਰਕ ਕਰੋ।

اگر آپ کو یہ معلومات کسی اور فارمیٹ یا زبان میں درکار ہوں، تو
برائے مہربانی مندرجہ بالا شعبے میں ہم سے رابطہ کریں.

"إذا احتجت الحصول على هذه المعلومة بشكل مغاير أو مترجمة إلى لغة مختلفة فيرجى منك الاتصال بالقسم المذكور أعلاه"